

# Higher order topological invariants from the Chern-Simons action

Roman V. Buniy<sup>1,\*</sup> and Thomas W. Kephart<sup>2,†</sup>

<sup>1</sup>*Institute of Theoretical Science, University of Oregon, Eugene, OR 94703*

<sup>2</sup>*Department of Physics and Astronomy,  
Vanderbilt University, Nashville, TN 37235*

(Dated: February 2, 2008)

It is well known that for a field theory with the Chern-Simons action, expectation values of Wilson line operators are topological invariants. The standard result is expressed in terms of the Gaussian linkings of closed curves defining the operators. We show how judicious choice of Wilson lines leads to higher order topological linkings.

## I. INTRODUCTION

Our understanding of knots and links in topological field theory began with the work of Polyakov [1] who showed that the expectation value of a Wilson line in the Chern-Simons theory gives the Gaussian linking of the components of a link. This work was soon followed by Witten's [2] extensive exploration of expectation values of Wilson lines in the non-abelian Chern-Simons theory which he showed is a natural framework for understanding Jones polynomials from the knot theory. Perturbative expansion of the expectation values leads to the third order linking and the corresponding link polynomials [5]. Our approach is somewhat different. Wilson loops in the abelian Chern-Simons theory also implicitly contains the requisite apparatus for the study of higher order linking, and we show here that this can be facilitated by a set of gauges [3, 4] tailored to the higher order linking problem. We begin with a discussion of intersection number and its relation to Gaussian linking. Next we review the minimal required background from Poincaré duality and the de Rham theorem needed to carry out our calculations. We then introduce a special set of gauge potentials that allow us to arrive at our main result—higher order linking invariants at all orders.

## II. INTERSECTION AND LINKING NUMBERS

Suppose  $C$  and  $C'$  are disjoint oriented closed curves in  $\mathbb{R}^3$ , and  $S$  and  $S'$  are surfaces such that  $\partial S = C$  and  $\partial S' = C'$ . By smooth deformations of  $C$  which leave it in  $\mathbb{R}^3 - C'$ , we can change the number of points of intersection of  $C$  and  $S'$ . To find a quantity which is invariant under such deformations, we note that additional points come in pairs, and intersections of  $C$  and  $S'$  have opposite orientations for points in each pair. If  $C$  and  $S'$  intersect transversely at a point  $p$ , we define the intersection index  $I(C, S', p)$  to be equal 1 or  $-1$  depending on the relative orientation of  $C$  and  $S'$  at  $p$ . We also define the intersection

---

\*Electronic address: roman@uoregon.edu

†Electronic address: tom.kephart@gmail.com

number  $I(C, S')$  as the sum of the intersection indices over all points of the intersection,

$$I(C, S') = \sum_{p \in C \cap S'} I(C, S', p). \quad (1)$$

It is clear that  $I(C, S')$  is invariant under the above deformations as contributions due to additional pairs of points cancel.

One can also deform  $S'$  into  $\tilde{S}'$  and notice that  $I(C, S') = I(C, \tilde{S}')$ , where  $\partial S' = \partial \tilde{S}' = C'$ . This means that the intersection number depends only on  $C$  and  $C'$ ; for this reason we call it the linking number,  $L(C, C') = I(C, S')$ . Examination of relevant orientations gives  $L(C, C') = L(C', C)$ . The method of Green functions leads to an explicit expression for the linking number of the curves in terms of their parametrizations,

$$L(C, C') = (4\pi)^{-1} \int_{C \times C'} \sum_{abc} \varepsilon_{abc} \frac{(x - x')^a}{\|x - x'\|^3} dx^b \wedge dx'^c. \quad (2)$$

### III. DUALITY

From the Poincaré duality and de Rham theorem [6, 7], for a closed curve  $C$  in three dimensions, there exists a closed 2-form  $F$  such that for any 1-form  $B$  we have

$$\int_C B = \int_{\mathbb{R}^3} B \wedge F. \quad (3)$$

We call  $(C, F)$  a dual set. Since  $B$  is arbitrary, it is clear that  $\text{supp } F = C$ . Since  $\mathbb{R}^3$  is simply connected, there exists a 1-form  $A$  such that  $F = dA$ . The Stokes theorem then gives [8]

$$\int_S dB = \int_{\mathbb{R}^3} dB \wedge A. \quad (4)$$

This means that there is a particular solution  $A$  such that  $\text{supp } A = S$ ,

$$A(x) = \int_{y \in S} \delta(x - y) \sum_a dx^a * dy^a. \quad (5)$$

We can similarly introduce a dual set  $(C', F')$  such that  $F' = dA'$  and for any 1-form  $B'$  we have

$$\int_{C'} B' = \int_{\mathbb{R}^3} B' \wedge F'. \quad (6)$$

Taking  $B = A'$ ,  $B' = A$  and using the Stokes theorem again, we find

$$\int_C A' = \int_{C'} A = \int_{\mathbb{R}^3} A \wedge dA' = L(C, C'). \quad (7)$$

Since  $L(C, C')$  is a topological invariant, smooth deformations of  $C$  which leave it in  $\mathbb{R}^3 - C'$  should not change its value. This is possible if and only if  $dA'|_{\mathbb{R}^3 - C'} = 0$ . If  $A'$  is exact, then  $L(C, C') = 0$ , and so a nontrivial case is when  $A'$  is a closed 1-form which is not exact.

#### IV. SECOND ORDER FIELDS

Suppose  $\{C_i\}_{1 \leq i \leq N}$  are disjoint closed curves in  $\mathbb{R}^3$  and let  $F_i = dA_i$  be dual to  $C_i$ . We now construct a dual set  $(C_{ij}, F_{ij})$  of the second order which satisfies

$$\int_{C_{ij}} B = \int_{\mathbb{R}^3} B \wedge F_{ij} \quad (8)$$

for any 1-form  $B$  and  $i \neq j$ . Since  $B$  is arbitrary, it follows that  $\text{supp } F_{ij} = C_{ij}$ . The most general 2-form which can be expressed in terms of  $A_i$  and  $A_j$  is

$$F_{ij} = f_i dA_j - f_j dA_i + g A_i \wedge A_j, \quad (9)$$

where  $f_i$ ,  $f_j$  and  $g$  are arbitrary functions. In order for  $F_{ij}$  to be closed, the functions have to satisfy certain conditions. A requirement  $dF_{ij}|_{\mathbb{R}^3 - C_i - C_j} = 0$  gives  $dg|_{\mathbb{R}^3 - C_i - C_j} = 0$  and without loss of generality we set  $g = 1$ . Requirements  $dF_{ij}|_{C_i} = 0$  and  $dF_{ij}|_{C_j} = 0$  then give

$$(df_j - A_j)|_{C_i} = 0, \quad (10)$$

$$(df_i - A_i)|_{C_j} = 0. \quad (11)$$

Integrating these conditions, we find a constraint  $L(C_i, C_j) = 0$ . This means that the second order field associated with a pair of closed curves can be defined only if the curves are unlinked.

Since  $dF_{ij} = 0$ , there exists a 1-form  $A_{ij}$  such that  $F_{ij} = dA_{ij}$ . We seek a solution in the form

$$A_{ij} = \frac{1}{2}(\gamma_i A_j - \gamma_j A_i). \quad (12)$$

From a requirement  $F_{ij} = dA_{ij}$ , the unknown functions  $\gamma_i$  and  $\gamma_j$  are found to satisfy

$$(d\gamma_i - A_i)|_{\mathbb{R}^3 - C_i - C_j} = 0, \quad (13)$$

$$d\gamma_i|_{C_i} = 0, \quad (14)$$

$$(d\gamma_i - 2A_i)|_{C_j} = 0, \quad (15)$$

$$(2f_i - \gamma_i)|_{C_j} = 0 \quad (16)$$

and the same expressions with  $i$  and  $j$  interchanged. This implies  $\gamma_i|_{\mathbb{R}^3 - \cup_i C_i} = \int_{\Gamma_i} A_i$ , where  $\Gamma_i$  is a curve in  $\mathbb{R}^3 - \cup_i C_i$ ; this means  $\gamma_i$  is a nonlocal quantity. If  $S_i \cap C_j = \emptyset$ , then  $d\gamma_i|_{C_j} = 0$ , and so  $\gamma_i|_{C_j}$  is a constant. If  $S_i \cap C_j \neq \emptyset$ , then  $S_i \cap S_j = \cup_m (S_i \cap S_j)_{(m)}$ , where  $m$  labels disjoint segments of the intersection; for an example, see Fig. 1. Let  $C_{j(m)}$  be the segment of  $C_j$  which closes the curve  $(S_i \cap S_j)_{(m)}$  and agrees with its orientation; this closed curve is  $C_{ij(m)} = C_{j(m)} \cup (S_i \cap S_j)_{(m)}$ . We define  $C'_j = \cup_m C_{j(m)}$  and its complement in  $C_j$  is  $C''_j$ . It follows from the above relations that  $\gamma_i|_{C'_j}$  and  $\gamma_i|_{C''_j}$  are constants such that  $\gamma_i|_{C'_j} - \gamma_i|_{C''_j} = 2$ . Without loss of generality, we set  $\gamma_i|_{C'_j} = 2$  and  $\gamma_i|_{C''_j} = 0$ . Using the definition of  $F_{ij}$ , the duality condition now becomes

$$\int_{C_{ij}} B = - \int_{C'_i} B + \int_{C'_j} B + \int_{S_i \cap S_j} B. \quad (17)$$




FIG. 1: Two components of  $C_{ij}$  are drawn with thick lines.  $S_i$  is a disk bounded by  $C_i$ , and the thick line segments in this disk are two disjoint segments of  $S_i \cap S_j$ .

This implies  $C_{ij} = C_i'^{-1} \cup C_j' \cup (S_i \cap S_j)$ , which agrees with  $\text{supp } F_{ij} = C_{ij}$ . See Fig. 1 for an example of the above construction. The Stokes theorem gives

$$\int_{S_{ij}} dA = \int_M dA \wedge A_{ij}, \quad (18)$$

where a surface  $S_{ij}$  is such that  $\partial S_{ij} = C_{ij}$ . This means  $\text{supp } A_{ij} = S_{ij}$ .

The above construction requires  $L(C_i, C_j) = 0$ . Since the curves  $C_i$  and  $C_j$  are unlinked, the surfaces  $S_i$  and  $S_j$  can be chosen to be disjoint, in which case  $dA_{ij}|_{\mathbb{R}^3 - (C_i \cup C_j)} = 0$ . One can similarly proceed to construct higher order fields. For example,  $A_{ijk}$  for  $i \neq j \neq k$  is required to satisfy  $dA_{ijk}|_{\mathbb{R}^3 - (C_i \cup C_j \cup C_k)} = 0$ . If all linkings of order  $q$ , where  $1 \leq q \leq p \leq N-1$  vanish, then we can similarly construct dual sets  $(C_{I_q}, dA_{I_q})$ , where  $I_q = (i_1, \dots, i_q)$  and  $i_1 \neq i_2 \neq \dots \neq i_q$ . The quantities  $dA_{I_q}$  are related to what is known in algebraic topology as the Massey products of cohomology groups [9]; see also [10].

## V. PATH INTEGRAL

We want to construct a field theory for which expectation values of observables are topological invariants of various orders. For this we need to specify the action and the observables. We choose the Chern-Simons action,

$$S(B) = \int_{\mathbb{R}^3} B \wedge dB, \quad (19)$$

since it has all the necessary topological properties [2]. First, it does not depend on the choice of metric. Second, it can be related to linking numbers in the following way. Suppose  $\Gamma_\alpha$  is a closed curve on which  $dB$  takes a constant value  $dB_\alpha$ . (In order for  $\Gamma_\alpha$  to be closed, we may need to identify points at infinity by considering  $S^3$  instead of  $\mathbb{R}^3$ . We also avoid field configurations with sources, like in monopoles.) Since a union of curves  $\Gamma = \cup_\alpha \Gamma_\alpha$  densely fills  $\mathbb{R}^3$ , an arbitrary 1-form  $B$  can be written as  $B = \sum_\alpha B_\alpha$ . The action becomes

$$S(B) = \sum_{\alpha\beta} L(\Gamma_\alpha, \Gamma_\beta), \quad (20)$$

and so (apart from the choice of measure for  $\Gamma$ ) we interpret it as the self-linking  $L(\Gamma, \Gamma)$  of the set of closed field lines of  $dB$ .

If  $C = \cup_{\alpha} C_{\alpha}$  is a union of disjoint closed curves, then an integral  $\int_C B$  is invariant with respect to deformations of  $C$  if and only if  $dB = 0$ . Since  $dB = 0$  is the classical equation of motion for  $S(B)$ , the integral is a topological observable (at least in the semiclassical limit). Proceeding as above, we find

$$\int_C B = \sum_{\alpha\beta} L(\Gamma_{\alpha}, C_{\beta}), \quad (21)$$

which we interpret as the linking  $L(\Gamma, C)$  of the sets of closed field lines of  $dB$  with  $C$ . Since the measure in the path integral is  $\exp(iS(B))$ , it is convenient to consider as an observable a Wilson loop operator  $W(C, B) = \exp(i \int_C B)$ .

We thus need to compute the expectation value

$$Z(C) = \int DB \exp(iS(B)) W(C, B). \quad (22)$$

Using duality, this becomes

$$Z(C) = \int DB \exp\left(i \int_{\mathbb{R}^3} B \wedge d(B + A)\right), \quad (23)$$

where  $dA$  is dual to  $C$ . Changing the variable  $B = B' - \frac{1}{2}A$  gives four terms in the exponent. One corresponds to the path integral of the measure without a Wilson loop,  $Z(\emptyset)$ , the two mixed terms combine into a boundary term, and the forth is the linking invariant. Ignoring the boundary term, we find

$$Z(C) = Z(\emptyset) \exp\left(-\frac{1}{4}i(L(C, C))\right). \quad (24)$$

If we take  $C = \cup_p \cup_{I_p} C_{I_p}^{n_{I_p}}$ , where  $C_{I_p}^{n_{I_p}}$  is a curve  $C_{I_p}$  repeated  $n_{I_p} \in \mathbb{Z}$  times, then we find

$$L(C, C) = \sum_{pq} \sum_{I_p J_q} n_{I_p} n_{J_q} L(C_{I_p}, C_{J_q}). \quad (25)$$

Here  $L(C_{I_p}, C_{J_q})$  is the first order linking of curves  $C_{I_p}$  and  $C_{J_q}$ ; it can also be interpreted as a linking of order  $p + q - 1$  of the curves  $\{C_i\}_{1 \leq i \leq N}$ . In general,  $L(C, C)$  is a sum of all linkings of orders  $1 \leq r \leq 2N - 1$ . We now look at two simple examples.

## VI. EXAMPLES

The simplest example is when only the second order linkings appear. We take  $C = \cup_{1 \leq i \leq N} C_i$  and find

$$L(C, C) = \sum_{ij} L(C_i, C_j), \quad (26)$$

in agreement with a result in Refs. [1, 2]. For the third order linking, we consider the simplest example [11, 12] of the Borromean rings in Fig. 2. Since  $L(C_i, C_j) = 0$  for  $i \neq j$ , we can




FIG. 2: The Borromean rings.

define the corresponding  $C_{ij}$ , and so we take  $C = C_1 \cup C_2 \cup C_3 \cup C_{12} \cup C_{23} \cup C_{31}$ . The second order linking vanishes and the third order linking is given by  $L_3(C, C) = \sum_{ijk} \varepsilon_{ijk} L(C_{ij}, C_k)$ . Comparison of Figs. 1 and 2 shows that  $L(C_{12}, C_3) = 1$ . This can also be seen from

$$L(C_{12}, C_3) = \int_{\mathbb{R}^3} A_1 \wedge A_2 \wedge A_3 - \frac{1}{2} \int_{C_1} \gamma_2 A_3 + \frac{1}{2} \int_{C_2} \gamma_1 A_3 \quad (27)$$

since in the special gauge the first term equals 1, two contributions to the second term cancel, and the third term vanishes. The symmetry then leads to  $L_3(C, C) = 6$ .

## VII. CONCLUSIONS

In the Chern-Simons theory, the first order (Gaussian) linking of two curves  $C_1$  and  $C_2$  associated with the 1-forms  $A_1$  and  $A_2$  can be ascribed to the topological properties of the expectation value that the associated Wilson lines are found to obey. Here we have generalized the idea to the case of higher order linking. First, if  $C_1$  and  $C_2$  are unlinked, then we are free to define a new second order 1-form  $A_{12}$  via Eq. (11). The associated curve  $C_{12}$ , unlike  $C_1$  and  $C_2$ , is not fixed in space, but is however sufficient for our topological needs, which are to investigate its linking with other curves. If for instance we have a third curve  $C_3$  unlinked with both  $C_1$  and  $C_2$ , but linked with  $C_{12}$ , then we find nonzero second order linking, i.e., a nonvanishing second order topological invariant, and a detailed treatment of the Borromean rings we given as an example of our method. The argument generalizes to higher orders. For curves  $\{C_i\}_{1 \leq i \leq N}$ , with no pairwise linking, no tertiary linking, etc., only  $(N - 1)$ th order linking, the general results are obtained from the path integral of the expectation value of  $N$  Wilson lines in the Chern-Simons gauge theory. One could apply these results to the investigation of higher order link polynomials.

### Acknowledgments

The work of RVB was supported by DOE grant number DE-FG06-85ER40224 and that of TWK by DOE grant number DE-FG05-85ER40226.

- 
- [1] A. M. Polyakov, *Mod. Phys. Lett. A* **3**, 325 (1988).
  - [2] E. Witten, *Commun. Math. Phys.* **121**, 351 (1989).
  - [3] R. V. Buniy and T. W. Kephart, arXiv:hep-th/0611334.
  - [4] R. V. Buniy and T. W. Kephart, arXiv:hep-th/0611335.
  - [5] E. Guadagnini, M. Martellini and M. Mintchev, *Phys. Lett. B* **227** (1989) 111; P. Cotta-Ramusino, E. Guadagnini, M. Martellini and M. Mintchev, *Nucl. Phys. B* **330** (1990) 557; E. Guadagnini, M. Martellini and M. Mintchev, *Nucl. Phys. B* **330** (1990) 575.
  - [6] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley-Interscience, New York, 1978.
  - [7] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
  - [8] We assume that the value of appropriate quantities vanishes at infinity, so there is no contribution from the boundary term. Boundary terms also vanish if we use  $S^3$  instead of  $\mathbb{R}^3$ .
  - [9] W. S. Massey, *Symp. Int. Topologia Algebraica*, Mexico, 145 (1959); W. S. Massey, *Proc. Conf. on Algebraic Topology*, Chicago, University of Illinois at Chicago, p. 174 (1968); D. Kraines, *Trans. Amer. Math. Soc.* **124**, 431 (1966); E. J. O'Neill *Trans. Amer. Math. Soc.* **248**, 37 (1979); R. A. Fenn, *Techniques of Geometric Topology*, Cambridge University Press, Cambridge, 1983; M. I. Monastyrsky and V. S. Retakh, *Commun. Math. Phys.* **103**, 445 (1986).
  - [10] M. A. Berger, *J. Phys. A*, **23**, 2787 (1990).
  - [11] D. Rolfsen, *Knots and Links*, Publish or Parish, Houston, 1990.
  - [12] L. Kauffman, *Knots and Physics*, World Scientific, Singapore, 2001.